arXiv:0811.4466v1 [quant-ph] 27 Nov 2008

Conservation rules for entanglement transfer between qubits

Stanley Chan,[1] M. D. Reid,[2] and Z. Ficek[1]
[1] Department of Physics, School of Physical Sciences,
The University of Queensland, Brisbane, QLD 4072, Australia
[2] ARC Centre of Excellence for Quantum-Atom Optics, School of Physical Sciences,
The University of Queensland, Brisbane QLD 4072, Australia

We consider an entangled but non-interacting qubit pair a_1 and b_1, where a_1 and b_1 are each independently coupled to a set of local qubit systems, a_I and b_J, respectively. We derive two general rules to describe the transfer of entanglement from a_1-b_1 to a_I-b_J, for the case where the initial entanglement is given by the two different Bell states, Ψ and Φ, corresponding to anti-correlated and correlated qubits respectively. We find that for Ψ, the non-local pairwise entanglement conserves the sum of the square of the non-local pairwise concurrences (SSPC). For Φ, the SSPC is reduced, even to zero in some cases, to reveal a complete and abrupt loss of all non-local pairwise entanglement.

Entanglement is not only crucial to the transition between classical and quantum behaviour [1], but is essential to many key applications in quantum information [2]. An understanding of how entanglement is transferred between systems and the existence of associated conservation rules is of fundamental and practical importance.

The seminal works of Bell and Einstein, Podolsky and Rosen (EPR) focused on entanglement that is shared between two distant and non-interacting "qubit" (two-level) systems [3]. This "nonlocal" entanglement can be maintained over long distances with exciting implications for both tests of quantum mechanics and applications such as quantum cryptography [4]. However, a fundamental issue is the degradation of entanglement brought about because each party inevitably interacts "locally" with other systems, as illustrated in Figure 1. This local coupling can lead to an abrupt depletion of all the original entanglement [5], an effect which has been recently experimentally confirmed [6]. The dynamical behaviour of entanglement under the action of the environment is regarded as a central issue in quantum information [7, 8].

Intuition tells us that the two-qubit entanglement is not truly "lost", but is simply redistributed among the interacting parties [8, 9, 10]. While it is the case that entanglement will be created between local systems, due to the local couplings, we would also expect a conservation of a global nonlocal entanglement, to reflect that no further "nonlocal" interaction has taken place.

This raises the natural question of whether a universal rule exists, for arbitrary local interactions, to describe the entanglement transfer among the qubits. More than this, there is the intriguing issue of how to relate the nonlocal entanglement between constituents of the two nonlocal systems (Fig. 1), with the global nonlocal entanglement that would be conserved.

Yonac *et al.* [9] have presented a conservation rule for the transfer of entanglement from one entangled qubit pair, a_1, b_1, into another, a_2, b_2. They observed $C_{11}+C_{22}$ to be constant, where C_{IJ} is the concurrence measure [12] of entanglement between qubits a_I and b_J. However, this simple conservation rule only applies to certain types of entanglement, that of Bell states $|\Psi\rangle$ with anti-correlated qubits, and is not universal to all local interactions. Lopez *et al.* [8] presented a very different result, for Bell states $|\Phi\rangle$ with correlated qubits. They showed that when such a qubit pair is coupled to independent reservoirs, the death of qubit entanglement precedes the birth of reservoir entanglement, so there is a temporary loss of all nonlocal pairwise entanglement. Again, their results applied only to certain local interactions.



Figure 1: We derive rules to describe how the entanglement is transferred among qubits, when each member of an entangled non-interacting qubit pair a_1, b_1 is locally and independently coupled to a set of qubits, a_I and b_I, respectively.

In this paper, we give new insight by deriving conservation rules for entanglement transfer between qubits that apply generally to all cases of "local" interactions that can exist within the qubit sets, $A \equiv \{a_I\}$ and $B \equiv \{b_J\}$. We consider the transfer of entanglement from one qubit pair a_1, b_1 into a set of qubits pairs a_I, b_J, $(1 < I \leq N, 1 < J \leq M)$, where members of all "non-local" pairs a_I, b_J are non-interacting (Fig 1). Which conservation rule is valid depends only on the type of global entanglement that is imprinted onto the system. Surprisingly, when the global entanglement is that of the Bell state $|\Psi\rangle$ with anti-correlated qubits, a remarkedly simple conservation result exists:

$$\sum_{I=1}^{N}\sum_{J=1}^{M} C_{IJ}^2(t) = C_{AB}^2. \tag{1}$$

The total nonlocal pairwise entanglement is conserved, but this manifests in the square of the concurrence, not in the concurrence itself**.** The conserved quantity is equal to the initial entanglement C^2_{AB}, which we show is also that of the global two-qubit system $\{A, B\}$. For this case, we will prove an additivity of constituent entanglement, that the entanglement shared between any two nonlocal partitions is the sum of the nonlocal pairwise entanglement of the constituents of the partitions. This rule applies to closed systems, and is investigated for open systems.

Where the global entanglement is that of Bell state $|\Phi\rangle$, the inequality

$$0 \leq \sum_{I,J} C^2_{IJ}(t) \leq C^2_{AB}. \tag{2}$$

holds. We will show there can be a vanishing of all nonlocal pairwise constituent entanglement C_{IJ}, despite conservation of global nonlocal entanglement C_{AB}, to give consistency with the result of Lopez *et al.* [8].

The two different scenarios may be thought of in the following way. Suppose entanglement exists between A and B, where A is made up of subsystems measured by Alice, Ann, Agatha respectively, and B is composed of subsystems measured by Bob, Bill and Brian. In the Ψ scenario, the global nonlocal entanglement can *always* be evaluated, through measurements shared only by pairs: Alice-Bob, Alice-Bill, Ann-Bob, Ann-Bill, etc. In the Φ scenario, this is not the case. Examples exist where all pairwise entanglement would be zero, despite there being a global entanglement, between A and B. This has potential implications for quantum cryptography, where measurement of shared entanglement between two parties, A and B, at different locations is used to determine security [13]. We will show however, that in both cases, the global entanglement could be inferred with communication between all parties at A and one at B: Alice-Ann-Agatha-Bob; Alice-Ann-Agatha-Bill, and so on.

We begin by considering the two non-interacting qubit systems, which we label A and B and which have outcomes 0 and 1 for some system observable Q_A and Q_B, respectively. Eigenstates are $|i\rangle|j\rangle$ where $i, j = 0, 1$ are outcomes for Q_A, Q_B respectively. Of interest is the case where the qubit represents one or zero energy quanta. We examine cases where each system A, B has constituent subsystems, and there are internal interactions between them, described by Hamiltonians H_A and H_B:

$$H = H_A + H_B. \tag{3}$$

A measure of bipartite entanglement between two qubit systems, such as A and B, is the concurrence, defined as $C_{AB} = \max\left(0, \sqrt{\lambda_1} - \sqrt{\lambda_2} - \sqrt{\lambda_3} - \sqrt{\lambda_4}\right)$, where λ_i are the eigenvalues of the density matrix $\rho' = \rho(\sigma_y \otimes \sigma_y)\rho^*(\sigma_y \otimes \sigma_y)$, where ρ is the system density operator and σ_y is the Pauli matrix in y direction. The maximum possible entanglement is given by $C_{AB} = 1$, while $C_{AB} = 0$ implies no entanglement. The state is entangled when $C_{AB} > 0$.

The global bipartite entanglement defined for systems A and B must be invariant throughout the evolution. This is understood since the Hamiltonian Eq. (3) does not allow interaction, or exchange of quanta, between the two systems, or with external systems. The evolution of the pure state $|\psi(0)\rangle = \sum_{i,j} c_{ij}|i\rangle|j\rangle$ is given by $|\psi(t)\rangle = e^{-iHt/\hbar}|\psi(0)\rangle = |\psi(0)\rangle$, since H_A, H_B conserve the bit values of A and B, respectively. The same argument applies if we begin with a mixture of pure states. Hence, the concurrence C_{AB}, defined from ρ, is invariant.

We now examine two special cases of pure state entanglement that can characterize the two qubit system $A-B$ [5]. These correspond to the two classses of Bell states for the two qubit system. The original Bell-like state Ψ with anticorrelated "spins" is

$$|\Psi\rangle = \cos\alpha|1\rangle|0\rangle + e^{i\beta}\sin\alpha|0\rangle|1\rangle \tag{4}$$

while that of correlated "spins" is written

$$|\Phi\rangle = \cos\alpha|1\rangle|1\rangle + e^{i\beta}\sin\alpha|0\rangle|0\rangle. \tag{5}$$

In both cases the concurrence is $C_{AB} = 2\sin\alpha\cos\alpha$, to indicate maximal entanglement at $\alpha = \pi/4$.



Figure 2: Evolution of the two-qubit concurrences for different nonlocal partitions, for state $|\Psi\rangle$ with Hamiltonian Eq. (9). C^2_{11} (solid line); C^2_{22} (dashed line); $C^2_{12} = C^2_{21}$ (dash-dotted line); C^2_{AB} (thin solid line). The following rule, universal for $|\Psi\rangle$, holds: $SSPC = C^2_{AB}$. Here $\Delta_a = \Delta_b = 0$, $g_a = g_b$and $\alpha = \pi/6$. In this case of symmetric interactions $g_a = g_b$, the simple conservation rule $C_{AB} = C_{11} + C_{22}$ of Yonac *et al.* [9] also holds.

Next we examine what happens when each of the systems A and B is composed of a collection of *interacting* qubits, which we denote $\{a_I\}$, $\{b_J\}$ (Fig. 1). Since the overall qubit value at each of A and B must be conserved under action of Eq. (3), for states $|\Psi\rangle, |\Phi\rangle$ there can be at most one qubit at each of A or B with a bit 1; all other bits are 0. Possible states are written $|\{1\}_n\rangle|\{1\}_m\rangle$ where $|\{1\}_n\rangle = |0...\{1\}_n..0\rangle$ denotes that the qubit a_n is in state $|1\rangle$ while all other qubits $a_{I\neq n}$ are in $|0\rangle$; $|\{1\}_m\rangle$ defines the qubit states of B. The state with zero for all qubits is written $|0\rangle$.

The evolution when the global entanglement is of form $|\Psi\rangle$ can be written $|\Psi(t)\rangle = \sum_{I=1}^{N} d_{AI}(t)|\{1\}_I\rangle|0\rangle + \sum_{J=1}^{M} d_{BJ}(t)|0\rangle|\{1\}_J\rangle$, where the $d_{AI}(t)$, $d_{BJ}(t)$ satisfy

$$\sum_{I=1,N} |d_{AI}(t)|^2 = \cos^2\alpha\,, \quad \sum_{J=1,M} |d_{BJ}(t)|^2 = \sin^2\alpha, \quad (6)$$

This conservation of probability holds because there is no external coupling, so that the total energy is conserved [10]. We now relate the global bipartite entanglement C_{AB}, which is conserved, to the non-local pairwise concurrences C_{IJ} of subsystems a_I and b_J. This pairwise concurrence is calculated by tracing over all other systems. Calculation shows the reduced density matrix for the a_I, b_J system to be of "X-state" form [9] (we omit the explicit time dependence in coefficients): we have

$$\rho_{IJ} = \begin{pmatrix} |d_{AI}|^2 & d_{AI}d_{BJ}^* & 0 \\ d_{AI}^* d_{BJ} & |d_{BJ}|^2 & 0 \\ 0 & 0 & \sum_{i\neq I}|d_{Ai}|^2 + \sum_{j\neq J}|d_{Bj}|^2 \end{pmatrix}, \quad (7)$$

in terms of the a_I, b_J basis states $|10\rangle$, $|01\rangle$ and $|00\rangle$. The pairwise concurrence is $C_{IJ} = 2|d_{AI}d_{BJ}|$.

Theorem: For the case of a global Bell entanglement C_{AB} of type $|\Psi\rangle$ between two non-interacting systems A and B, the sum of the square of the pairwise constituent "nonlocal" concurrences (SSPC) is conserved: $SSPC = \sum_{I=1}^{N}\sum_{J=1}^{M} C_{IJ}^2 = C_{AB}^2$. The entanglement shared by *any* two nonlocal partitions $\{a_i, a_j, ...\}$, $\{b_m, b_n...\}$ satisifes a simple Pythagorean addition of constituent entanglement:

$$C^2_{\{i,j,...\}\{m,n,...\}} = \sum_{k=i,j,..}\ \sum_{l=m,n,...} C^2_{kl}. \quad (8)$$



Figure 3: Evolution of two- qubit concurrences for different nonlocal partitions, for state $|\Psi\rangle$ with Hamiltonian Eq. (9). The plots confirm the rule, universal for $|\Psi\rangle$, that $SSPC = C_{AB}^2$. C_{11} (solid line); C_{22} (dashed line); C_{12} (dashed-dotted line); C_{21} (dotted line). Here $\Delta_a = \Delta_b = 0$, $g_a = 2g_b$, $g = (g_a + g_b)/2$ and $\alpha = \pi/4$.

Proof: For any Hamiltonian of the form $H = H_A + H_B$, the probability sums, Eq. (6), are constant [10]. Hence, given that the pairwise concurrence is derived from Eq. (7), the conjecture must hold: $SSPC = \sum_I\sum_J |C_{IJ}|^2 = 4(\sum_{I=1,N}|d_{AI}(t)|^2) \times (\sum_{J=1,M}|d_{BJ}(t)|^2) = 4\cos^2\alpha\sin^2\alpha = C_{AB}^2$. The result Eq. (8) follows in the same manner from direct evaluation of concurrences after tracing.

We give a verification of the conservation rule by examining the case of $N, M = 2$ where the qubits a_1, b_1 are two-level atoms with transition frequency ω_0 and the qubits a_2, b_2 are cavity modes with resonant frequencies ω_a, ω_b. We consider where the local interaction Hamiltonian is of the Jaynes-Cummings form [11]:

$$H_A = \hbar\omega_0 S_A^z + \hbar\omega_a\left(a^\dagger a + \frac{1}{2}\right) + \hbar g_a\left(a^\dagger S_A^- + aS_A^\dagger\right). \quad (9)$$

Here S_A^+, S_A^- and S_A^z are raising, lowering and spin-z operators for the atom qubit a_1, and $a^\dagger$ (a) are the creation (annihilation) operators for the cavity mode qubit a_2. The g_a is the strength of the coupling between the atom and the cavity mode. The local Hamiltonian for B is defined similarly. Solutions for the concurrences describing pairwise entanglement can be evaluated for the full case of arbitrary couplings and detunings, defined as $\Delta_a = (\omega_0 - \omega_a)/2$, $\Delta_b = (\omega_0 - \omega_b)/2$. Figures 2 and 3 show concurrences for this case, to confirm the conservation law, Eq. (1).

Where the global entanglement of the qubits a_1, b_1 is that of the Bell state $|\Phi\rangle$, the Pythagorean sum of the pairwise concurrences is no longer generally conserved, but satisfies the inequality Eq. (2). In this case, the wave function evolves as $|\Phi(t)\rangle = \sum_{i=1}^{N}\sum_{j=1}^{M} c_{i,j}(t)|\{1\}_i\rangle|\{1\}_j\rangle + c_0|0\rangle|0\rangle$. Here

$$|c_0(t)|^2 = \sin^2\alpha, \quad \sum_{i,j}|c_{ij}(t)|^2 = \cos^2\alpha \quad (10)$$

to describe that the quanta at A and B are shared among the local qubits. In this case the reduced density matrix ρ_{IJ} for the qubits a_I and b_J is, in terms of basis states $|11\rangle$, $|10\rangle$, $|01\rangle$ and $|00\rangle$,

$$\begin{pmatrix} |c_{IJ}|^2 & 0 & 0 & c_{IJ}c_0^* \\ 0 & \sum_{n\neq J}|c_{In}|^2 & 0 & 0 \\ 0 & 0 & \sum_{m\neq I}|c_{mJ}|^2 & 0 \\ c_{IJ}^* c_0 & 0 & 0 & \sum_{n,m\neq J,I}|c_{m,n}|^2 + |c_0|^2 \end{pmatrix}. \quad (11)$$

The concurrence is $C_{IJ} = \max\left\{0, 2(|c_{IJ}||c_0| - \sqrt{\left(\sum_{n\neq J}|c_{In}|^2\right)\left(\sum_{m\neq I}|c_{mJ}|^2\right)})\right\}$. We note immediately that $C_{IJ}^2 \leq 4|c_0|^2|c_{IJ}|^2$ and thus, using Eq. (10), the inequality Eq. (2) must hold. In fact, we show below that all nonlocal pairwise entanglement C_{IJ} can vanish ($SSPC = 0$), even where maximal global entanglement

C_{AB} exists. We note however this is not the case for all nonlocal partitions. Simple evaluation reveals for $|\Phi\rangle$ states that entanglement C_{AJ} defined for nonlocal partitions $\{a_1, ..., a_N\}\,\{b_J\}$ satisfies a simple addition rule: $\sum_J C_{AJ}^2 = C_{AB}^2$.

The loss of all nonlocal pairwise concurrence for evolution of the Bell state $|\Phi\rangle$ is evident in the model Eq. (9) (Fig. 4). Where the entanglement C_{AB} is low, we can identify regions where each C_{IJ} is zero (SSPC= 0). The SSPC is regained when the transfer of entanglement fom qubits a_1, b_1 into the qubits a_2, b_2 is complete. We observe, in this case of low C_{AB}, the feature observed by Lopez *et al.* [8] for reservoir interactions,that the "birth" of entanglement in cavity modes is delayed a finite time after the "death" of entanglement in the atoms.

As a final note we show that the constituent addition rule Eq. (1) for $|\Psi\rangle$ also applies to describe entanglement transfer for open systems, where energy loss is modelled by zero-temperature reservoir interactions: $H_R = a_I\Gamma^\dagger + a_I^\dagger\Gamma$ if the qubits are cavity modes, or $H_R = \sigma_I\Gamma^\dagger + \sigma_I^\dagger\Gamma$ if the qubits are two-level atoms. Here $\Gamma = \sum_R g_R b_R$ and b_R is the boson destruction operator for one of many vacuum modes that comprise the reservoir. Inclusion in the Hamiltonian Eq. (9) of such coupling, under the Markovian assumption and assuming equal cavity and atomic damping rates γ, leads to a simple damping envelope for the concurrences: $C_{IJ}(t) \rightarrow C_{IJ}(t)\exp(-\gamma t)$. The global entanglement C_{AB} shows asymptotic decay $C_{AB}(t) = C_{AB}(0)\exp(-\kappa t)$, where κ is the global decay rate, to confirm the rule $\sum_{IJ} C_{IJ}^2(t) = C_{AB}^2$ (Fig. 5).

This work was funded by the Australian Research Council and the ARC Center of Excellence program.



Figure 4: Evolution of pairwise concurrences for different nonlocal partitions, for state $|\Phi\rangle$ and Hamiltonian Eq. (9). Here $\Delta_a = \Delta_b = 0$ and $g_a = g_b = g$. The top figure plots SSPC for $\alpha = \pi/4$ (solid line), $\alpha = \pi/6$ (dashed line), $\alpha = \pi/12$ (dashed-dotted line). The bottom figure plots the individual concurrences for $\alpha = \pi/12$: C_{11}^2 (solid line), C_{22}^2 (dashed line), C_{AB}^2 (thin solid line). $C_{12}^2 = C_{21}^2$ (dash-dotted line) is below 0.01in this case. As the global entanglement C_{AB} weakens, regions of total loss of nonlocal pairwise entanglement are evident (SSPC=0).

---

[1] A. Einstein, B. Podolsky and N. Rosen, Phys. Rev. **47**, 777 (1935); J. S. Bell, Physics **1,** 195 (1964); A. Peres, Phys. Rev. Lett. **76**, 1413 (1996).
[2] M. A. Nielsen and I. L. Chuang, Quantum Computation and Quantum Information (Cambridge University Press. Cambridge, 2000).
[3] A. Zeilinger, Rev. Mod. Phys. **71**, S288 (1999).
[4] R. Ursin et al, Nature Physics, **3**, 483 (2007).
[5] T. Yu and J. H. Eberly, Phys. Rev. Lett. **93**, 140404 (2004); L. Diosi, Lect. Notes Phys. **622**, 157 (2003); P. J. Dodd and J. J. Halliwell, Phys. Rev. A 69, 052105 (2004); Z. Ficek and R. Tanas, Phys. Rev. A **74**, 024304 (2006); M. F. Santos, P. Milman, L. Davidovich and N. Zagury, Phys. Rev. A **73**, 040305 (2006); Anna Jamroz, J. Phys. A **39**, 7727 (2006); M. O. Terra Cunha, New J. Phys. **9**, 237 (2007); A. Al-Qasimi and D. F. V. James, Phys. Rev. A **77**, 012117 (2008); L. Aolita, R. Chaves, D. Cavalcanti, A. Acin, and L. Davidovich, Phys. Rev. Lett. **100**, 080501 (2008).



Figure 5: Evolution of pairwise concurrences for state $|\Psi\rangle$ and Hamiltonian Eq. (9) with coupling to reservoirs. C_{11}^2 (dashed line); C_{22}^2 (dashed-dotted line); $C_{12}^2 = C_{21}^2$ (dotted line); C_{AB}^2 (solid line). Here $\Delta_a = \Delta_b = 2$, $g_a = g_b$, $\alpha = \pi/4$, $\gamma = 0.3$.

[6] M. P. Almeida et al., Science **316**, 579 (2007).

[7] J. H. Eberly and T. Yu, Science **316** (5824), 555 (2007).

[8] C. E. Lopez, G. Romero, F. Lastra et al, Phys. Rev. Lett. **101**, 080503 (2008).

[9] M. Yonac, T. Yu and J. H. Eberly, J. Phys. B, At Mol Opt Phys. **39** S621 (2006); M. Yonac, T. Yu, and J. H. Eberly, J. Phys. B, At Mol Opt Phys. **40**, S45 (2007).

[10] I. Sainz and G. Bjork, Phys. Rev. A **76**, 042313, (2007).

[11] E. T. Jaynes and F. W. Cummings, Proc. *IEEE* **51** 89 (1963).

[12] K. Wootters, Phys. Rev. Lett. **80**, 2248 (1998).

[13] A. Ekert, Phys. Rev. Lett. **67**, 661 (1991).